


QA
343
W6

UC-NRLF



5C 164 718



UNIVERSITY OF
CALIFORNIA

THE ELLIPTIC MODULAR FUNCTIONS
ASSOCIATED WITH THE ELLIPTIC
NORM CURVE E^7

BY

ROSCOE WOODS

A. B. GEORGETOWN COLLEGE, 1914

A. M. UNIVERSITY OF MAINE, 1916

OCT 30 1920

THESIS

SUBMITTED IN PARTIAL FULFILLMENT OF THE REQUIREMENTS FOR THE

DEGREE OF

DOCTOR OF PHILOSOPHY

IN MATHEMATICS

IN

THE GRADUATE SCHOOL

OF THE

UNIVERSITY OF ILLINOIS

1920

Reprinted from the TRANSACTIONS OF THE AMERICAN MATHEMATICAL SOCIETY,
Vol. 23, No. 2, March, 1922.

70 7000
ABSTRACT

UNIV. OF
CALIFORNIA

THE ELLIPTIC MODULAR FUNCTIONS
ASSOCIATED WITH THE ELLIPTIC
NORM CURVE E^7

BY

ROSCOE WOODS

A. B. GEORGETOWN COLLEGE, 1914

A. M. UNIVERSITY OF MAINE, 1916

7.
1/2

THESIS

SUBMITTED IN PARTIAL FULFILLMENT OF THE REQUIREMENTS FOR THE

DEGREE OF

DOCTOR OF PHILOSOPHY

IN MATHEMATICS

IN

THE GRADUATE SCHOOL

OF THE

UNIVERSITY OF ILLINOIS

1920

Reprinted from the TRANSACTIONS OF THE AMERICAN MATHEMATICAL SOCIETY,
Vol. 23, No. 2, March, 1922.

70 .vml
ABPOMIAO

1343
-rfg

2

ACKNOWLEDGMENT

To Professor A. B. Coble under whose supervision this paper has been written I am especially indebted for his very valuable suggestions, his deep interest and his unfailing kindness and encouragement.

UNIVERSITY OF ILLINOIS, MAY, 1920.

ROSCOE WOODS.

CONTENTS

	Page
Introduction.....	179
I. The groups connected with E^7	
1. The group $G_{2,7^2}$ of collineations of the E^7 into itself.....	180
2. The fixed heptahedra of the 8 cyclic G_7 's.....	181
3. A canonical form of the $G_{2,7^2}$	182
4. The family of E^7 's.....	182
5. The fixed spaces.....	184
II. The quadrics on E^7	
1. The pencil of quadrics on E^7	185
2. The group on the quadrics.....	186
3. A Kleinian form.....	187
III. The interpretation of the form F'	
1. Its fundamental elliptic modular functions.....	187
2. The null-system.....	188
3. The rational curves in S_2 and S_3	189
4. The net of quadrics in $S_3(\alpha)$	190
5. The modular line and spread.....	191
IV. The loci in S_3	
1. The net of quadrics in S_2	193
2. The plane of the half period points.....	194
3. The locus of the zero point in S_3	195
4. Summary.....	196

THE ELLIPTIC MODULAR FUNCTIONS ASSOCIATED WITH THE ELLIPTIC NORM CURVE E^n *

BY

ROSCOE WOODS

INTRODUCTION

The elliptic norm curve E^n in space S_{n-1} admits a group G_{2n^2} of collineations and there is a single infinity of such curves which admit the same group. A particular E^n of the family is distinguished by the coördinates of a point on a modular curve, the ratios of these coördinates being elliptic modular functions defined by the modular group congruent to identity (mod n). In the group G_{2n^2} there are certain involutory collineations with two fixed spaces. If E^n is projected from one fixed space upon the other, a family of rational curves C^m mapping the family of E^n 's is obtained. The quadratic irrationality separating involutory pairs on E^n involves the coördinates of a point on the modular curve and the parameter t on a member of the family C^m .

Miss B. I. Miller† has discussed the elliptic norm curves for which $n = 3, 4, 5$. In these cases the genus of the modular group is zero and a point of the modular curve can be denoted by a value of the binary parameter τ . The irrationality separating involutory pairs on E^n was used by her to define an elliptic parameter

$$u = \int \frac{(tdt)}{\sqrt{(t\tau)\alpha_\tau^{r-3}\alpha_t^3}}$$

where α_τ^r is the tetrahedral, octahedral, or icosahedral form. This form of u is invariant under all the cogredient transformations of t and τ which leave α_τ^r unaltered.

The cases considered by Dr. Miller are relatively simple, due to the fact that the genus of the modular group is zero. In this paper, the case $n = 7$ for which the genus is 3, one which is fairly typical of the general case, is subjected to a

* Presented to the Society, April 14, 1922.

† See these Transactions, vol. 17 (1916), p. 259.

similar investigation. Many of the results may be extended to the case where n is any prime number and in some features to the case where n is any odd number. By methods of geometry and group theory, we derive in this discussion the well known elliptic modular functions attached to this group as well as some new ones and obtain a number of their algebraic properties.* This treatment suggests a number of "root functions," i. e., square roots of modular functions which are themselves uniform.

In §I, the groups and subgroups associated with the E^7 are discussed and thrown into a canonical form. The equations of the transformation from S_6 to the fixed spaces S_2, S_3 , and the equations of the groups of transformations in these spaces are derived. These have been found without the aid of function theory and have been checked with Klein's results in Klein-Fricke's *Elliptische Modulfunktionen*. In §II, a single Kleinian form† is derived which furnishes the fourteen linearly independent quadrics whose complete intersection is E^7 . From this form in §III the fundamental elliptic modular functions $t_1 : t_2 : t_4$ are determined. Also the families C^2, C^3 of rational curves in S_2 and S_3 are found. In §IV, the loci in S_3 are discussed. The paper closes with a parametric representation of E^7 .

I. THE GROUPS CONNECTED WITH E^7

1. **The group $G_{2,7^2}$ of collineations of E^7 into itself.** The homogeneous coördinates of a point of the elliptic norm curve E^7 are $x_0 : x_1 : \dots : x_6 = 1 : p(u) : p'(u) : \dots : p''(u)$. As u runs over the period parallelogram ω_1, ω_2 the E^7 is obtained in a six-dimensional space S_6 . It is known‡ that the only birational transformations of the general elliptic curve into itself are given by $u' = \pm u + b$, where b is any constant.§ From the parametric representation of the E^7 as set forth above, it is evident that seven points of the E^7 on a hyperplane section are characterized by the fact that the sum of their parameters is congruent to zero (mod ω_1, ω_2) and conversely. In view of this, all transformations for which $7b \equiv 0 \pmod{\omega_1, \omega_2}$ are collineations. This congruence has three irreducible solutions

$$(1) \quad b = 0, \quad b = \omega_1/7, \quad b = \omega_2/7.$$

* In the case $n = 4$, Miss Miller has expressed the opinion that the properties of the elliptic integral associated with E^4 and the Dyck quartic should apply to Klein's quartic which occurs in this case. This has not been verified.

† By a Kleinian form is meant a form in several variables invariant under isomorphic linear groups on these variables.

‡ Appell-Goursat, *Fonctions Algébriques*, p. 474.

§ Segre, *Mathematische Annalen*, vol. 27(1887), p. 296.

Klein-Fricke, *Theorie der elliptischen Modulfunktionen*, vol. 2, p. 241. Hereafter the initials K. F. will be used to refer to this work.

These furnish the substitutions

$$(2) \quad \begin{aligned} S_{01} : u' &= u + \omega_{01}, \\ S_{10} : u' &= u + \omega_{10}, & \omega_{ij} &= i\omega_1/7 + j\omega_2/7, \\ V : u' &= -u, & & (i, j = 0, 1, \dots, 6). \end{aligned}$$

S_{01} and S_{10} are collineations of period seven and generate a group G_7 , which is abelian in its elements. V is a collineation of period two which adjoined to G_7 , generates a group $G_{2 \cdot 7}$. This group $G_{2 \cdot 7}$ of collineations contains all the collineations of the general E^7 into itself.

The G_7 in the $G_{2 \cdot 7}$ contains 8 cyclic G_7 's and no other subgroups. These are denoted by $G_\infty, G_1, \dots, G_6$ where G_∞ is generated by S_{01} and G_i by $S_{10}S_{01}^i$ ($i = 0, 1, \dots, 6$). The elements of $G_{2 \cdot 7}$ not in G_7 are of the form

$$(3) \quad V_{ij} : u' = -u + \omega_{ij} \quad (i, j = 0, 1, \dots, 6),$$

and are of period two. The V_{ij} form a conjugate set. Any cyclic G_7 with one involution generates a dihedral $G_{2 \cdot 7}$ which contains seven involutions. Hence there are 56 dihedral G 's. These with the cyclic G_7 's complete the subgroups of $G_{2 \cdot 7}$. The relations satisfied by the generators of $G_{2 \cdot 7}$ are

$$(4) \quad \begin{aligned} S_{01}^7 &= S_{10}^7 = 1, \\ S_{01}S_{10} &= S_{10}S_{01}, \quad VS_{10} = S_{10}^{-1}V, \quad VS_{01} = S_{01}^{-1}V. \end{aligned}$$

2. **The fixed heptahedra of the 8 cyclic G_7 's.** The condition that a hyperplane section touch the E^7 in seven coincident points is given by

$$(5) \quad 7u \equiv 0 \pmod{\omega_1, \omega_2}.$$

The irreducible solutions of this congruence furnish the 49 parameters ω_{ij} of the singular points. Under G_∞ the 49 points ω_{ij} separate into 7 sets of seven conjugate points such that each set is on a hyperplane. Such a set of seven hyperplanes will be called a *heptahedron*. Since there are 8 cyclic G_7 's, there are 8 heptahedra which will be designated by $H_\infty, H_0, \dots, H_6$.*

The 49 singular points are now arranged in a matrix (using only the subscripts) in such a way that the rows furnish the 7 sets of conjugate points which determine the 7 hyperplanes of H_∞ , while the columns furnish the 7 hyperplanes of H_0 .

* The reason for calling one heptahedron H_x will appear later. These heptahedra can be determined from the resolvent equation of the 8th degree associated with the Galois problem of degree 168. Compare K. F., vol. 1, p. 732.

$$(6) \quad \left\| \begin{array}{cccccc} 00 & 01 & 02 & 03 & 04 & 05 & 06 \\ 10 & 11 & 12 & 13 & 14 & 15 & 16 \\ 20 & 21 & 22 & 23 & 24 & 25 & 26 \\ 30 & 31 & 32 & 33 & 34 & 35 & 36 \\ 40 & 41 & 42 & 43 & 44 & 45 & 46 \\ 50 & 51 & 52 & 53 & 54 & 55 & 56 \\ 60 & 61 & 62 & 63 & 64 & 65 & 66 \end{array} \right\|$$

Each row of this matrix is transformed into itself by G_∞ , each column into itself by G_0 . Further the seven hyperplanes of each heptahedron are linearly independent. Let us prove this for G_∞ . If the seven hyperplanes X_i are not independent, there is a relation among them involving $7 - k$ of these X 's such that these $7 - k$ X 's do not satisfy further relations. Then the $7 - k$ X 's in this relation are all fixed under G_∞ and meet in an S_k which is also fixed under G_∞ . Hence G_∞ permutes the S_5 's on the fixed S_k in such a way that $7 - k$ of the S_5 's are fixed. Therefore by projection from S_k upon an S_{5-k} , we should have in S_{5-k} $7 - k$ fixed spaces no. $6 - k$ of which were related. But such a collineation is the identity in S_{5-k} . Hence every S_5 on S_k is fixed, contrary to the fact that G_∞ has only a finite number 7 of fixed spaces.

3. **A canonical form of the $G_{2,7}$.** Let then the heptahedron H_∞ with linearly independent faces be chosen as a reference figure and denote these faces by X_i ($i = 0, 1, \dots, 6$).^{*} These are determined by the rows of the matrix (6). The reference figure is completed by choosing a unit-hyperplane. This hyperplane will be chosen as the one containing the singular points of the first column of the matrix (6).

In terms of the coördinates thus defined the generators of the $G_{2,7}$ of collineations of the E^7 into itself have the form

$$(7) \quad \begin{array}{l} S_{01} : X'_i = X_{i-1} \quad (X_{7+i} \equiv X_i) \\ S_{10} : X'_i = \epsilon^i X_i \quad (i = 0, 1, \dots, 6) \\ V : X'_i = X_{-i} \quad (X_{-i} \equiv X_{7-i}) \end{array}$$

where ϵ is a seventh root of unity. The formulas (7) constitute a first canonical form of $G_{2,7}$.

4. **The family of E^7 's.** The curve E^7 depends upon the ratio $\omega = \omega_1/\omega_2$. For each value of ω , there is an E^7 , hence there is a family F of E^7 's. But the

^{*} X_i is written instead of $X_i(u)$. The X_i can be represented as the products of sigma functions, i. e.,

$$X_i(u) = \alpha_i \prod_{j=0}^6 \sigma(u - \omega_j)$$

where the α_i are constants which insure the double periodicity of the ratios X_i . Compare K. F., vol. 2, p. 238.

group $G_{2,7}$ of collineations is the same for each member of the family F since its coefficients are numbers independent of the ratio ω . For each curve of F the set of 8 heptahedra is the same, since the heptahedra are determined by their common $G_{2,7}$.

All collineations which leave each member of F unaltered have been determined. If there are further collineations which interchange the members of F , they must arise from integer period transformations of determinant $+1$. Consider then the transformation

$$(8) \quad \begin{aligned} \omega_1' &= \alpha\omega_1 + \beta\omega_2 & \alpha\delta - \beta\gamma &= 1 \\ \omega_2' &= \gamma\omega_1 + \delta\omega_2 \end{aligned}$$

where $\alpha, \beta, \gamma, \delta$ are integers. The curve as first expressed in terms of $p(u)$ and its derivatives is unaltered by (8). In the new reference system the curve and each ω_{ij} are unaltered if (8) is congruent to identity (mod 7). On the other hand if (8) is not congruent to identity (mod 7), the ω_{ij} are permuted and we may look upon this operation either as merely a change in the coördinate system in which the curve is fixed or as a collineation in which the reference system is fixed and the E^7 passes into a new curve which belongs to F . Therefore all transformations (8) which are congruent to identity (mod 7) give rise to the identical collineation. These transformations constitute a subgroup of (8) of index 2·168. All elements of (8) in a coset of this subgroup give rise to a collineation which permutes the curves of F except the element $\begin{vmatrix} -1 & 0 \\ 0 & -1 \end{vmatrix}$ which is the element V . Hence there are 2·168/2 or 168 collineations which interchange the members of F .* These collineations may be represented by the elements of (8) reduced modulo 7, that is

$$(9) \quad \begin{aligned} \omega_1' &= \alpha\omega_1 + \beta\omega_2 & \alpha\delta - \beta\gamma &\equiv 1 \pmod{7} \\ \omega_2' &= \gamma\omega_1 + \delta\omega_2 \end{aligned}$$

It is well known that any transformation of the group (8) is a combination of the transformations

$$(10) \quad S : \omega' = \omega + 1 \quad T : \omega' = -1/\omega,$$

where S is of period 7 and T is of period 2 when reduced modulo 7. Since $T^2 \equiv S^7 \equiv (ST)^3 \equiv (S^4T)^4 \equiv 1 \dagger$, these relations define a G_{168} of collineations on the reduced periods which permutes the members of the family F . Therefore we have the following theorem:

* See K. F., vol. 1, p. 398.

† It should be noted that in homogeneous form, T is of period 4, (S^4T) is of period 8. Hence V and T^2 are the same. Compare Dickson, *Linear Groups*, p. 303.

THEOREM I. *The family F of elliptic E^7 's, each member of which is unaltered by $G_{2,7^2}$, is unaltered as a whole by a collineation group $G_{2,7^2 \cdot 168}$ for which $G_{2,7^2}$ is an invariant subgroup. Under the group of F each curve belongs to a conjugate set of 168 curves.**

The collineation T permutes the H_i ($i = \infty, 0, 1, \dots, 6$) as follows: $(\infty 0)$, (16) , (25) , (34) , where the subscripts only are used. The collineation S permutes the H_i ($i = 0, 1, \dots, 6$) cyclically and leaves H_∞ invariant. Under the group (8) the H_i are permuted like the 8 points $\infty, 0, 1, \dots, 6$ in a finite geometry modulo 7, there being 8 points on a line.

The equations of the collineations S and T in terms of X_i are†

$$(11) \quad \begin{aligned} S : X'_i &= \epsilon^{-i^2/2} X_i \\ T : X'_i &= c \sum_{\beta=0}^6 \epsilon^{i\beta} X_\beta \quad (i = 0, 1, \dots, 6). \end{aligned}$$

5. **The fixed spaces.** In $G_{2,7^2}$, the 7^2 involutions V_{ij} ($i, j = 0, 1, \dots, 6$) have the form $u' = -u + \omega_{ij}$. The fixed points of these involutions are $u \equiv \omega_{ij}/2 + P/2$ where $P/2$ can evidently have the values $0, \omega_1/2, \omega_2/2$, and $(\omega_1 + \omega_2)/2$. We consider the simplest set, i. e., the set for which $i = j = 0$.

Due to the involutory character of V , there are two skew spaces of fixed points in S_6 , an S_2 and an S_3 . If the coördinates of these fixed spaces be denoted by y_i and z_j ($i = 0, 1, 2, 4; j = 1, 2, 4$) respectively, the equations of the transformation from the coördinates X_i to those of y and z are

$$(12) \quad \begin{aligned} X_0 &= y_0, & X_1 - X_6 &= 2z_1, \\ X_1 + X_6 &= 2y_1, & X_2 - X_5 &= 2z_2, \\ X_2 + X_5 &= 2y_2, & X_4 - X_3 &= 2z_4, \\ X_4 + X_3 &= 2y_4, & & \end{aligned}$$

In terms of y and z , V now has the form

$$(13) \quad y'_i = y_i, \quad z'_j = -z_j \quad (i = 0, 1, 2, 4; j = 1, 2, 4).$$

In (12), $y_i = 0$ determine the S_2 of fixed points and $z_j = 0$ determine the S_3 of fixed points. The fixed S_5 's are either on S_2 with equations $\alpha_0 y_0 + \alpha_1 y_1 + \alpha_2 y_2 + \alpha_4 y_4 = 0$ or on S_3 with equations $\beta_1 z_1 + \beta_2 z_2 + \beta_4 z_4 = 0$. The α 's may be determined by putting the S_5 on u_1, u_2, u_3 , three arbitrary points on E^7 , so that necessarily this S_5 cuts E^7 in the points $-u_1, -u_2, -u_3$. Therefore the S_5 contains the point $u = 0$, but no proper half period point. Hence all the fixed S_5 's on the S_2 and therefore S_2 itself, contain the point $u = 0$ but no proper

* See K. F., vol. 1, p. 398.

† Compare K. F., vol. 2, p. 292. The formula for S is compatible with Klein's for n a prime number. As we deal with collineations in homogeneous forms we do not need to keep c of the K. F. formula; it is therefore dropped in the remainder of the work.

half period point. Therefore S_3 contains the proper half period points since they are also fixed points.

The family F of E^7 's projected from the fixed S_2 upon the fixed S_3 becomes a family F_1 of rational cubics doubly covered, since the pairs ($\neq u$) corresponding under V each project into the same point. In a similar manner, by projection from S_3 upon S_2 , F becomes a family F_2 of conics doubly covered.

It is my purpose to discuss the families F_1, F_2 , for which the curves in each family will vary with ω whereas the points on a particular curve will vary with the pairs ($\neq u$) on the original E^7 . The $G_{2.72.168}$ has now reduced to a G_{168} in S_2 and S_3 which leaves F_1 and F_2 invariant. This G_{168} is generated by S and T whose equations are easily found to be

$$(14) \quad \begin{aligned} S &: \begin{aligned} y'_i &= \epsilon^{-i/2} y_i \\ z'_j &= \epsilon^{-j/2} z_j \end{aligned} \quad (i = 0, 1, 2, 4; j = 1, 2, 4) \\ T &: \begin{aligned} y'_i &= y_0 + \sum_j (\epsilon^{ij} + \epsilon^{-ij}) y_j \quad (i = 0, 1, 2, 4) \\ z'_k &= \sum_l (\epsilon^{lk} - \epsilon^{-lk}) z_l \quad (j, k, l = 1, 2, 4). \end{aligned} \end{aligned}$$

Formulas (12), (13) and (14) constitute a second canonical system of coördinates for E^7 .

II. THE QUADRICS ON E^7

1. **The pencil of quadrics on E^7 .** Hermite has shown that the number of linearly independent quadrics on E^7 is fourteen. These fourteen quadrics cut out the E^7 completely with no extraneous intersection.* In the second system of coördinates a general quadric has the form

$$(15) \quad q_0 = \sum_{i, k=0}^6 \alpha_{ik} X_i X_k = 0,$$

where α_{ik} are constants. Let us suppose that the α_{ik} are so determined that the quadric contains the curve E^7 . Under the collineation S_{10} , E^7 is transformed into itself. Hence the quadric (15) is transformed into a quadric on E^7 . The transforms of q_0 under S_{10} are of the form

$$(16) \quad q_j = \sum_{i, k=0}^6 \alpha_{ik} \epsilon^{j(i+k)} X_i X_k = 0 \quad (j = 0, 1, \dots, 6).$$

Since each q_j is on E^7 , a linear combination of them will be on the curve. Multiplying each q_j by unity and adding we obtain a particular quadric Q_0 on E^7 characterized by the fact that it consists only of those terms for which $i + k \equiv 0 \pmod{7}$. Using the multipliers 1, ϵ^5 , ϵ^3 , ϵ^1 , ϵ^6 , ϵ^4 , ϵ^2 , respectively, we obtain a

* Compare K. F., vol. 2, p. 245.

second *particular* quadric Q_1 on E^7 characterized by the fact that it consists only of those terms for which $i + k \equiv 1 \pmod{7}$. Proceeding in this way we obtain 7 *particular* quadrics on E^7 . They are

$$(17) \quad Q_i = \alpha_{i0}X_i^2 + 2\alpha_{i1}X_{i+1}X_{i-1} + 2\alpha_{i2}X_{i+2}X_{i-2} + 2\alpha_{i4}X_{i+4}X_{i4} \\ = 0 \quad (i = 0, 1, \dots, 6).$$

Any quadric on the curve E^7 is a linear combination of the Q 's, since the seven Q 's contain as yet 28 arbitrary coefficients. But since each Q_i is sent into Q_{i+1} by S_{01} , these 28 coefficients reduce to four, i. e., $\alpha_0, \alpha_1, \alpha_2, \alpha_4$. From these seven Q 's, we know that we must be able to get the 14 linearly independent quadrics on the E^7 . The α 's therefore must contain a parameter linearly and there will be one quadric of the type Q_i for which a particular α will vanish.* At most, then, a pencil can arise from the four terms of each Q_i . Any one of these seven pencils is defined by the fact that it admits one of the seven dihedral $G_{2,7}$'s whose cyclic subgroup is S_{10} . For example Q_0 admits the dihedral $(S_{10}V)$.

Since the α 's contain a parameter linearly, they may be interpreted as the coordinates of a point on a line in an S_3 . By choosing two members from the pencil of quadrics, the line is determined. We shall determine the α 's later as functions of ω and the parameter just mentioned.

2. **The group on the quadrics.** Under $G_{2,7}$, each member of the family F of E^7 's is transformed into itself and the quadrics on each curve are transformed into quadrics on that curve, so that a group of collineations is induced upon the Q_i as variables. Moreover since S and T interchange the members of the 168 sets of conjugate curves, they will send the quadrics on a given curve into a linear combination of the quadrics on the transformed curve. If we indicate the group $G_{2,7 \cdot 168}$ on the X_i 's in (7) and (11) by $G(\epsilon)$, then the induced group on the quadrics Q_i is $G(\epsilon^2)$.

In order to express all the quadrics (17) by one equation, consider the general quadric obtained by taking a linear combination of them. Such a quadric has the form.

$$(18) \quad \sum_{i=0}^6 L_i Q_i = 0,$$

where the L_i are arbitrary constants. On a given curve of F determined by a proper set of values of α_i ($i = 0, 1, 2, 4$), the bilinear form (18) is an identity in L and u . If we require that this bilinear form be an invariant under $G(\epsilon^2)$, there will be a certain group induced upon the L_i as variables. This group on the variables L_i is $G(\epsilon^{-2})$.

* Compare K. F., vol. 2, p. 268. Klein obtained the quadrics on the elliptic curves from the three-term sigma relation.

3. **A Kleinian form.** Since the properties of the groups on the L_i and Q_i are the same as those on the X_i , we isolate one of the involutions in the L_i , Q_i groups, i. e., that one induced by V which was isolated in the X_i group. We introduce the variables v and u , ζ and ϑ with Q_i and L_i , respectively, as y and z were introduced with the X_i . The equations of the transformations from Q_i and L_i to v , u , ζ and ϑ can be written down as were those for y and z . After this change of variables, (18) has the form

$$\begin{aligned}
 (19) \quad F' = & \zeta_0[\alpha_0 y_0^2 + 2\alpha_1 y_1^2 + 2\alpha_2 y_2^2 + 2\alpha_4 y_4^2 - 2\alpha_1 z_1^2 - 2\alpha_2 z_2^2 - 2\alpha_4 z_4^2] \\
 & + 2\zeta_1[\alpha_0 y_1^2 + 2\alpha_1 y_0 y_2 + 2\alpha_2 y_1 y_4 + 2\alpha_4 y_2 y_4 + \alpha_0 z_1^2 + 2\alpha_2 z_1 z_4 - 2\alpha_4 z_2 z_4] \\
 & + 2\zeta_2[\alpha_0 y_2^2 + 2\alpha_1 y_1 y_4 + 2\alpha_2 y_0 y_4 + 2\alpha_4 y_1 y_2 - 2\alpha_1 z_1 z_4 + \alpha_0 z_2^2 + 2\alpha_4 z_1 z_2] \\
 & + 2\zeta_4[\alpha_0 y_4^2 + 2\alpha_1 y_2 y_4 + 2\alpha_2 y_1 y_2 + 2\alpha_4 y_0 y_1 + 2\alpha_1 z_2 z_4 - 2\alpha_2 z_1 z_2 + \alpha_0 z_4^2] \\
 & + 4\vartheta_1[\alpha_0 y_1 z_1 + \alpha_1 y_0 z_2 - \alpha_2(y_1 z_4 + y_4 z_1) + \alpha_4(y_2 z_4 - y_4 z_2)] \\
 & + 4\vartheta_2[\alpha_0 y_2 z_2 + \alpha_1(y_4 z_1 - y_1 z_4) + \alpha_2 y_0 z_4 - \alpha_4(y_1 z_2 + y_2 z_1)] \\
 & + 4\vartheta_4[\alpha_0 y_4 z_4 - \alpha_1(y_2 z_4 + y_4 z_2) + \alpha_2(y_1 z_2 - y_2 z_1) + \alpha_4 y_0 z_1] = 0.
 \end{aligned}$$

On E^7 the above form is an identity in ζ , ϑ and can be separated into seven parts. However we shall have occasion to separate it into two parts, P_1 and P_2 , such that the part P_1 contains the coefficients ζ and the part P_2 the coefficients ϑ . The part P_1 is partly symmetrical and partly alternating in the coefficients α and ζ , hence the ζ 's can be interpreted as the coördinates of a point on a line in an S_3 and are therefore cogredient to the α 's. Hence we may conclude this section with the theorem

THEOREM II. *F' is a Kleinian form which remains invariant under the simultaneous transformation by the isomorphic groups $M(\epsilon)$ of (14) on the variables y and z ; $M(\epsilon^{-2})$ on the variables ζ and α and ϑ . The form F' determines the curve E^7 uniquely when the modular functions α are properly given, i. e., subject to the relation which connects their ratios.*

III. THE INTERPRETATION OF THE FORM F'

1. **Its fundamental elliptic modular functions.** Each curve of the family F has on it the point whose parameter is $u = 0$. As $\omega = \omega_1/\omega_2$ varies this zero point generates a locus. It has already been pointed out that the zero point is in the space S_2 of fixed points, i. e., when $u = 0$ all the y 's vanish. Let $z_i = t_i$ ($i = 1, 2, 4$) for $u = 0$; then F' in (19) reduces to

$$\begin{aligned}
 (20) \quad & \zeta_0[0 - 2\alpha_1 t_1^2 - 2\alpha_2 t_2^2 - 2\alpha_4 t_4^2] \\
 & + 2\zeta_1[\alpha_0 t_1^2 - 0 + 2\alpha_2 t_1 t_4 - 2\alpha_4 t_2 t_4] \\
 & + 2\zeta_2[\alpha_0 t_2^2 - 2\alpha_1 t_1 t_4 - 0 + 2\alpha_4 t_1 t_2] \\
 & + 2\zeta_4[\alpha_0 t_4^2 + 2\alpha_1 t_2 t_4 - 2\alpha_2 t_1 t_2 - 0] \equiv 0.
 \end{aligned}$$

Since (20) is an identity in the ζ_i , their coefficients must vanish. These coefficients are linear in the α 's, all of which do not vanish simultaneously, therefore the determinant of the α 's must vanish. After removing numerical factors, we find a skew-symmetric determinant of even order. This determinant is a perfect square.* It furnishes in variables t_i Klein's quartic, which is denoted as follows:

$$(21) \quad K = t_1^3 t_2 + t_2^3 t_4 + t_4^3 t_1 = 0.$$

K is the equation of the locus of the zero point of the family of E^7 's and admits a group G_{168} of collineations into itself, cogredient to the group in z in (14). The ratios $t_1 : t_2 : t_4$ are the *fundamental elliptic modular functions of the form F'* . The expressions for these ratios as uniform functions of the modulus ω may be obtained by setting $u = 0$ in the expressions for the z 's in terms of u , ω_1 , ω_2 , as indicated.

Since the curve E^7 varies with ω , and since each E^7 possesses a zero point, i. e., a point t which is on K , it is clear that the variation of E^7 with ω may be imaged by the variation of t on K . We shall express other elliptic modular functions associated with the family of E^7 's in terms of the t_i .

2. **The null-system.** The form in (20) is a null-system, since it can be written in the form

$$(22) \quad (\alpha_0 \zeta_1)^2 t_1^2 + (\alpha_0 \zeta_2)^2 t_2^2 + (\alpha_0 \zeta_4)^2 t_4^2 + 2(\alpha_4 \zeta_2) t_1 t_2 + 2(\alpha_1 \zeta_4) t_2 t_4 + 2(\alpha_2 \zeta_1) t_4 t_1 = 0,$$

where $(\alpha_i \zeta_k) = \alpha_i \zeta_k - \alpha_k \zeta_i$. Since (20) vanishes independently of the ζ 's it represents a *singular null-system*.† Hence (22) is the equation of a line whose coördinates may be taken as

$$(23) \quad \begin{aligned} (\alpha_0 \zeta_1) &= + 2 t_1 t_2, & (\alpha_4 \zeta_2) &= t_1^2, \\ (\alpha_0 \zeta_2) &= + 2 t_2 t_4, & (\alpha_1 \zeta_4) &= t_2^2, \\ (\alpha_0 \zeta_4) &= + 2 t_4 t_1, & (\alpha_2 \zeta_1) &= t_4^2, \end{aligned}$$

where α is clearly a point on a line. Since the coördinates of the line of the α 's are functions of t , we shall call it the *modular line* and denote it by L_m . The intersection of the coördinate planes of the reference tetrahedron in the space of the α 's, an $S_3^{(\alpha)}$, with L_m furnishes four convenient sets of values of the α 's, which substituted in F' give rise to the 28 quadrics on E^7 , of which only 14 are linearly independent, since any two sets of the α 's are linear combinations of the remaining two sets. These sets of values are

* Burnside and Panton, *Theory of Equations*, vol. 2, p. 46.

† See Veblen and Young, *Projective Geometry*, vol. 1, p. 324.

$$(24) \quad \alpha_0 : \alpha_1 : \alpha_2 : \alpha_4 = \begin{cases} 0 : -2t_1t_2 : -2t_2t_4 : -2t_1t_4 \\ 2t_1t_2 : 0 : +t_4^2 : -t_2^2 \\ 2t_2t_4 : -t_4^2 : 0 : t_1^2 \\ 2t_1t_4 : t_2^2 : -t_1^2 : 0 \end{cases}$$

The sets (24) suggest that we make a transformation on the α 's in F' . Let ξ be a plane such that it intersects L_m in the point α . From (24) we find this transformation to be

$$(25) \quad \begin{aligned} \alpha_0 &= 0 + 2t_1t_2\xi_1 + 2t_2t_4\xi_2 + 2t_1t_4\xi_4, \\ \alpha_1 &= -2t_1t_2\xi_0 + 0 - t_4^2\xi_2 + t_2^2\xi_4, \\ \alpha_2 &= -2t_2t_4\xi_0 + t_4^2\xi_1 + 0 - t_1^2\xi_4, \\ \alpha_4 &= -2t_1t_4\xi_0 - t_2^2\xi_1 + t_1^2\xi_2 + 0. \end{aligned}$$

If F' is transformed by (25), it will take the form

$$(26) \quad \sum \xi_i \zeta_j \phi_{ij} + \sum \xi_i \vartheta_l \phi'_{il} = 0 \quad (i, j = 0, 1, 2, 4 ; l = 1, 2, 4).$$

The 28 quadrics on the curve E^7 , of which naturally only 14 are linearly independent, are found by equating to zero the coefficients of the terms $\xi_i \zeta_j$ and $\xi_i \vartheta_l$ respectively, i. e., the ϕ_{ij} and ϕ'_{il} . We shall have occasion to use all of these quadrics, but will refer to them briefly in the above notation.

3. **The rational curves in S_2 and S_3 .** We have seen that α and ζ are cogredient variables and that P_1 is partly alternating and partly symmetrical in α and ζ . We now rewrite P_1 so as to exhibit this property. It has the form

$$(27) \quad \alpha_0 \zeta_0 \gamma_0^2 + 4 \sum \alpha_1 \zeta_1 \gamma_0 \gamma_2 + 2 \sum (\alpha_0 \zeta_1 + \alpha_1 \zeta_0) \gamma_1^2 + 4 \sum (\alpha_2 \zeta_4 + \alpha_4 \zeta_2) \gamma_1 \gamma_2 + 2 \sum (\alpha_0 \zeta_1) z_1^2 + 4 \sum (\alpha_4 \zeta_2) z_1 z_2 = 0,$$

where Σ , unless otherwise denoted, refers to the cyclic advance of the subscripts 1, 2, 4. This form furnishes the means by which the projections of the family F of E^7 's upon the fixed spaces S_2 and S_3 are found. The second part P_2 , bilinear in y and z , does not enter in these projections, since it vanishes when either space is considered separately.

Since ζ is perfectly arbitrary, consider it on the modular line L_m . Now interchange α and ζ in (27). The new form is similar to the old except that the sign of each term in z is changed. Denote the transformed P_1 by P'_1 . Since P_1 in (27) is a quadric on E^7 and since we consider ζ on L_m , P'_1 is also a quadric on E^7 . Whence their sum $P_1 + P'_1$ and their difference $P_1 - P'_1$ are quadrics on E^7 . Consider the former;

$$(28) \quad \alpha_0 \zeta_0 \gamma_0^2 + 4 \sum \alpha_1 \zeta_1 \gamma_0 \gamma_2 + 2 \sum (\alpha_0 \zeta_1 + \alpha_1 \zeta_0) \gamma_1^2 + 4 \sum (\alpha_2 \zeta_4 + \alpha_4 \zeta_2) \gamma_1 \gamma_2 = 0.$$

The equation (28) for arbitrary α and ζ on L_m furnishes a system of quadrics in S_3 which intersect in a cubic curve. From the symmetry of α and ζ in (28), we lose no generality by setting $\alpha_i = \zeta_i$. We then have

$$(29) \quad \alpha_0^2 \gamma_0^2 + 4 \sum \alpha_1^2 \gamma_0 \gamma_2 + 4 \sum \alpha_0 \alpha_1 \gamma_1^2 + 8 \sum \alpha_2 \alpha_4 \gamma_1 \gamma_2 = 0.$$

Since α is linear in a parameter λ on L_m , (29) furnishes a system of quadrics quadratic in λ . The coefficients of this quadratic system of quadrics are functions of t , so that as t varies on K , we get a family F_1 of cubic curves C^3 in S_3 . Hence we may state the following theorem:

(30) THEOREM III. *The projection C^3 of the curve E^7 upon S_3 is the base curve of the quadratic system of quadrics (29).*

Consider now the difference $P_1 - P'_1$. This is a conic in S_2 . It has the form

$$(31) \quad \sum (\alpha_0 \zeta_1) z_1^2 + 2 \sum (\alpha_4 \zeta_2) z_1 z_2 = 0,$$

which from (23) may be written as follows:

$$(32) \quad \sum t_1 t_2 z_1^2 + \sum t_1^2 z_1 z_2 = 0.$$

This shows that the system of conics varies with t on K . It is the polar conic of K as to z . Hence the theorem:

(33) THEOREM IV. *The projection C^2 of the family F of E^7 's upon S_2 is the system of polar conics of Klein's quartic K .*

4. The net of quadrics in S_3 (α). The quadric in (29) will be the square of a plane when the rank of its discriminant is 1. Its discriminant is of rank 1 if only the three relations

$$(34) \quad \alpha_0 \alpha_2 - \alpha_4^2 = 0, \quad \alpha_0 \alpha_4 - \alpha_1^2 = 0, \quad \alpha_0 \alpha_1 - \alpha_2^2 = 0,$$

are satisfied.

Consider now the net of quadrics

$$(35) \quad t_1(\alpha_0 \alpha_2 - \alpha_4^2) + t_2(\alpha_0 \alpha_4 - \alpha_1^2) + t_4(\alpha_0 \alpha_1 - \alpha_2^2) = 0.$$

From the transformations S and T in (14) we conclude that (35) is a Kleinian form.

The discriminant of the net (35) is K . Hence so long as t is on K , the quadric (35) has a double point. If we border the discriminant (35) with variables ξ and expand, we find the equation of this double point to be

$$(36) \quad (\alpha \xi)^2 = \xi_0^2 4 t_1 t_2 t_4 + \sum (-t_1^3 - t_2^2 t_4) \xi_1^2 + \sum (2 t_1 t_4^2) (2 \xi_0 \xi_1) + \sum (2 t_1^2 t_4) (2 \xi_1 \xi_2) = 0.$$

Therefore the coördinates of the double-point are

$$\begin{aligned}
 (37) \quad \rho \alpha_0 &= \sqrt{4t_1 t_2 t_4} & : 2t_1 t_2 t_4 & : t_1 t_4^2 & : t_1^2 t_2 & : t_2^2 t_4 \\
 \rho \alpha_1 &= \sqrt{-t_1^3 - t_2^2 t_4} & : t_1 t_4^2 & : -t_1^3 - t_2^2 t_4 & : t_1^2 t_4 & : t_4^2 t_2 \\
 \rho \alpha_2 &= \sqrt{-t_2^3 - t_4^2 t_1} & : t_2 t_1^2 & : t_1^2 t_4 & : -t_2^3 - t_4^2 t_1 & : t_2^2 t_1 \\
 \rho \alpha_4 &= \sqrt{-t_4^3 - t_1^2 t_2} & : t_4 t_2^2 & : t_4^2 t_2 & : t_2^2 t_1 & : -t_4^3 - t_1^2 t_2
 \end{aligned}$$

where ρ is 1, α_0 , α_1 , α_2 , α_4 , respectively. That is to say, we can express the entire system (37) rationally and without extraneous factors by giving the ten quadratic combinations of the α 's. These combinations are the coefficients of the terms ξ_i , ξ_k in (36).

The order of the linear modular group in the space of the y 's and α 's is double the order of the group* in the space of the z 's, that is, the group is a $G_{2,168}$, due to the fact that the identical collineation appears in the form $y_i = \epsilon y_i$. Hence the coördinates of a modular-point or plane in $S_3^{(\alpha)}$ and likewise in S_3 cannot be expressed rationally in terms of the t_i without an extraneous factor. The coördinates may however be expressed irrationally in terms of t as above, and it is to be noted that their ratios are uniform functions of ω .

A number of such modular root functions are suggested by the geometry of the system of cubic curves C^3 in S_3 . Thus the locus of the zero point on the curves C^3 , the locus of the plane of the half period points, the locus of the point where the tangent at the zero point meets the half period plane, as well as the transforms of these points and planes in the null-system of C^3 , give rise to functions of this type. Some of these are determined later.

The locus of the double point (36) as t varies on K is a well known space curve J of order 6 in $S_3^{(\alpha)}$,† whose points are in a one-to-one correspondence with the points of K . If we border the discriminant of (35) with ξ and η , which are to be thought of as parameters, we have ∞^6 curves of the third order in t which intersect K in 12 points which correspond to the 12 meets of the planes ξ , η with J . Hence when $\xi = \eta$ the cubic in t will be a contact cubic of K . Thus the system (36) for variable ξ is a system of contact curves of the third order associated with J .‡

5. **The modular line and spread.** If a point y be taken on J , a quadric of the net (35) has a node at y and the polar plane of this point as to this quadric vanishes, while the polar planes of the other two quadrics meet in a line. Take the coördinates of the point y on J as those in the second column of (37). The three

* Compare K. F., vol. 2, p. 313.

† Compare Snyder and Sisam, *Analytic Geometry of Space*, p. 168.

‡ See K. F., vol. 1, p. 716.

polar planes of this point as to the quadrics in the net (35) are in a pencil, and have the form

$$(38) \quad \begin{aligned} \alpha_0 t_1^2 - 0 + 2 \alpha_2 t_1 t_4 - 2 \alpha_4 t_2 t_4 &= 0, \\ \alpha_0 t_2^2 - 2 \alpha_1 t_1 t_4 - 0 + 2 \alpha_4 t_1 t_2 &= 0, \\ \alpha_0 t_4^2 + 2 \alpha_1 t_2 t_4 - 2 \alpha_2 t_1 t_2 - 0 &= 0. \end{aligned}$$

The axis of the pencil of planes (38) is the modular line L_m . Every point on L_m is in a one-to-one correspondence with the point y on J . Since the coördinates of L_m and of the point y on J are functions of t , the variation of y and of L_m also may be imaged by the variation of t on K . Hence as y generates J , L_m generates a ruled surface of order 8. That M is of order 8 may be shown as follows. The condition that a line l meet L_m is a linear condition on their coördinates, or a conic in t . This conic in t meets K in 8 points to each of which there corresponds a meet of l and M , whence M is of order 8.

Let us now consider the general quadric Q in the net (35), and put on it the condition that it have a node. The four partial derivatives $\partial Q / \partial \alpha_i$ must then vanish simultaneously. These are

$$(39) \quad \begin{aligned} 0 + \alpha_1 t_4 + \alpha_2 t_1 + \alpha_4 t_2 &= 0, \\ \alpha_0 t_4 - 2\alpha_1 t_2 &= 0, \\ \alpha_0 t_1 - 2\alpha_2 t_4 &= 0, \\ \alpha_0 t_2 - 2\alpha_4 t_1 &= 0. \end{aligned}$$

The discriminant of these equations is K . If we eliminate t from the equations (39), we find four cubic surfaces on each of which is J . Hence their common intersection is J . The equations of these are obtained from the vanishing of the third order determinants in the matrix of the equations (39). They are

$$(40) \quad \begin{aligned} S_1 &= \alpha_0^3 - 8 \alpha_1 \alpha_2 \alpha_4 = 0, \\ S_2 &= \alpha_0^2 \alpha_4 + 2 \alpha_0 \alpha_2^2 + 4 \alpha_1 \alpha_2^2 = 0, \\ S_3 &= \alpha_0^2 \alpha_1 + 2 \alpha_0 \alpha_4^2 + 4 \alpha_2 \alpha_4^2 = 0, \\ S_4 &= \alpha_0^2 \alpha_2 + 2 \alpha_0 \alpha_1^2 + 4 \alpha_4 \alpha_1^2 = 0. \end{aligned}$$

The modular spread M multiplied by α_0 is the following combination of S in (40):

$$(41) \quad S_1^3 - 8 S_2 S_3 S_4 = \alpha_0 M = 0.$$

From this result it is evident that J is a triple curve on M . Further, it can be shown that through every point of J there pass three trisecants of J and that L_m itself is a trisecant of J .*

This section can be partially summarized in the following theorem:

* The equation of M and the facts concerning J are easily obtained from a Cremona transformation of the third order.

THEOREM V. Through every point $\alpha (= \alpha_0, \alpha_1, \alpha_2, \alpha_4)$ on the octavic ruled surface M there passes a line L_m and the pencil of points α on L_m set in the form F' determines the quadrics on the curve E^7 . As the line L_m varies on M , the E^7 varies in the family F . The line L_m (itself a trisecant of J) meets the triple curve J on M in three points which correspond to the three trisecants of J that meet in a point t of J . Thus the points t of J are in a one-to-one correspondence with the curves of the family of E^7 's.

This completes the determination of the coefficients α of the quadrics F' which define the curve E^7 .

IV. THE LOCI IN S_3

1. The net of quadrics in S_3 . In (35) a net of quadrics in $S_3^{(\alpha)}$ was considered. The modular line L_m and the modular spread were associated with this net. Consider now a similar net of quadrics in plane coördinates U in S_3 , and let us find the condition that this net have a double plane. From the contragredient transformations S and T on the y 's in (14)*, we conclude that the following net is a Kleinian form:

$$(42) \quad t_1(2 U_0 U_1 - U_2^2) + t_2(2 U_0 U_2 - U_4^2) + t_4(2 U_0 U_4 - U_1^2) = 0.$$

The discriminant of this net is K . The bordered form of the discriminant is the square of a plane in point coördinates, i. e.,

$$(43) \quad t_1 t_2 t_4 y_0^2 + \sum (-t_2^3 - t_1 t_4^2) y_1^2 + 2 \sum t_1^2 t_2 y_0 y_1 + 2 \sum t_1 t_2^2 y_1 y_2 = 0.$$

So long as t is on K , the coördinates of the double plane (43) are

$$(44) \quad \begin{aligned} \rho U_0 &= \sqrt{t_1 t_2 t_4} & : & t_1 t_2 t_4 & : & t_1^2 t_2 & : & t_2^2 t_4 & : & t_4^2 t_1, \\ \rho U_1 &= \sqrt{-t_2^3 - t_1 t_4^2} & : & t_1^2 t_2 & : & -t_2^3 - t_1 t_4^2 & : & t_1 t_2^2 & : & t_1^2 t_4, \\ \rho U_2 &= \sqrt{-t_4^3 - t_2 t_1^2} & : & t_2^2 t_4 & : & t_1 t_2^2 & : & -t_4^3 - t_2 t_1^2 & : & t_2 t_4^2, \\ \rho U_4 &= \sqrt{-t_1^3 - t_4 t_2^2} & : & t_4^2 t_1 & : & t_1^2 t_4 & : & t_2 t_4^2 & : & -t_1^3 - t_4 t_2^2, \end{aligned}$$

where ρ is 1, U_0, U_1, U_2, U_4 , respectively. As in (37), we may express the entire system in (44) by taking the 10 quadratic combinations of the U 's from (43). The remarks following (37) apply here. The plane coördinates U_i taken from the second column of (44) are the modular systems A_γ developed by Klein. †

With the net (42) there will be a modular line L'_m , four cubic surfaces S'_i , a modular surface M' and a sextic J' . The coördinates of L'_m can be developed

* See K. F., vol. 1, p. 719.

† See K. F., vol. 1, p. 719.

in a manner similar to that used in finding those of L_m as the axis of the pencil of planes (38). They are

$$(45) \quad \begin{aligned} (U_0U'_1) &= t_1t_2, & (U_2U'_1) &= t_1^2, \\ (U_0U'_2) &= t_2t_4, & (U_4U'_2) &= t_2^2, \\ (U_0U'_4) &= t_4t_1, & (U_1U'_4) &= t_4^2. \end{aligned}$$

To every position of the plane $(Uy) = 0$ in (43) we have a line L'_m whose coördinates are given in (45). Since the coefficients of the plane $(Uy) = 0$ and L'_m are functions of t , the variation of the plane (Uy) and L'_m also may be imaged as the variation of t on K . It should be noted that the space of the α 's is different from the space of the y 's. Hence the modular lines L_m, L'_m ; the curves J, J' ; the spreads M, M' ; and the cubic surfaces S_i, S'_i are all distinct.

2. **The plane of the half period points.** For the three half period points, the z 's all vanish. If in the 14 linearly independent quadrics on E^7 we set the z 's all zero, we then obtain 8 quadrics in y (since 6 of the 14 quadrics are bilinear in y and z and vanish for z set equal zero). These 8 quadrics must pass through the half period points. If we call the plane of these points (Uy) , then we should be able to obtain from these 8 quadrics the four combinations $y_i(Uy)$ ($i = 0, 1, 2, 4$). The combinations furnishing these types of quadrics come from the systems

$$(46)^* \quad \begin{aligned} t_1t_4^3\phi_{12} + t_1t_2^2t_4\phi_{14} - t_1^2t_2^2\phi_{24} - t_2^2t_4^2\phi_{21} + t_2t_4^3\phi_{40} &= 0, \\ t_1\phi_{41} + t_2\phi_{01} - t_4\phi_{04} &= 0, \\ t_2\phi_{12} + t_4\phi_{02} - t_1\phi_{01} &= 0, \\ t_4\phi_{24} + t_1\phi_{04} - t_2\phi_{02} &= 0. \end{aligned}$$

The common factor (Uy) obtained from these equations (46) when the z 's are zero is precisely the plane

$$(47) \quad (Uy) = t_1t_2t_4ty_0 + t_1^2t_2y_1 + t_2^2t_4y_2 + t_4^2t_1y_4 = 0,$$

whose square appeared in (43). Hence the coördinates of the half period plane are the modular functions set forth in (44).

Since the half period plane is of the form $\Sigma \alpha_i y_i = 0$ ($i = 0, 1, 2, 4$), and since it may be considered as an S_5 in S_6 , it contains the point $u = 0$ and three pairs of points ($\pm u$) on E^7 , since the three pairs are sufficient to determine the α 's. It is therefore a fixed S_5 on the fixed S_2 . Since the pairs ($\pm u$) are the half period points, they are coincident points in S_6 , hence the half period plane (47) considered as an S_5 is a tritangent hyperplane of E^7 , tangent at the points $\omega_1/2, \omega_2/2$ and $(\omega_1 + \omega_2)/2$ and passing through the point $u = 0$.

* We draw from the entire system of quadrics ϕ_{ij} (16 in number) for convenience. These ϕ_{ij} are the coefficients of the terms $\xi_i \xi_j$ in F' after the transformation in (25).

Let us now consider the systems of quadrics in (46) with the z 's different from zero. These expressed in terms of y and z are

$$\begin{aligned}
 (48) \quad t_4 y_0(Uy) &= + 2t_1 t_2^2 z_2 (41) + (- 2t_1 t_2 t_4 z_2 - 2t_2^2 t_4 z_1) (12), \\
 t_4 y_1(Uy) &= - t_2^2 t_4 z_4 (12) + (t_2^2 t_4 z_2 - t_1^2 t_2 z_1) (41), \\
 t_1 y_2(Uy) &= - t_4^2 t_1 z_1 (24) + (t_4^2 t_1 z_4 - t_2^2 t_4 z_2) (12), \\
 t_2 y_4(Uy) &= - t_1^2 t_2 z_2 (41) + (t_1^2 t_2 z_1 - t_4^2 t_1 z_4) (24),
 \end{aligned}$$

where $(ik) = t_i z_k - t_k z_i$. Each of the above quadrics vanishes for $z_i = t_i$, that is each conic on the right in (48) intersects the polar conic C^2 in the zero point. The three remaining variable intersections of these conics and the polar conic correspond to the intersection of the plane $(\alpha_i y_i) = 0$ and the curve C^3 in S_3 . Hence the system of quadrics (48) give a parametric representation of the curve C^3 . To each z in (48) there is a definite point y in S_3 except at the base point of the system $z_i = t_i$. This representation can be put in a simpler form if we multiply the quadrics in (48) by $t_i t_k$, so that each quadric on the left has the common factor $t_1 t_2 t_4(Uy)$, which may be dropped, leaving the parametric representation of the curve C^3 as follows:

$$\begin{aligned}
 (49) \quad y_0 &= \rho [+ 2t_1^2 t_2^3 z_2 (41) + (- 2 t_1^2 t_2^2 t_4 z_2 - 2t_1 t_2^3 t_4 z_1) (12)], \\
 y_1 &= \rho [- t_1 t_2^3 t_4 z_4 (12) + (t_1 t_2^3 t_4 z_2 - t_1^3 t_2^2 z_1) (41)], \\
 y_2 &= \rho [- t_2 t_4^3 t_1 z_1 (24) + (t_2 t_4^3 t_1 z_4 - t_2^3 t_4^2 z_2) (12)], \\
 y_4 &= \rho [- t_4 t_1^3 t_2 z_2 (41) + (t_4 t_1^3 t_2 z_1 - t_4^3 t_1^2 z_4) (24)].
 \end{aligned}$$

Hence the doubled C^2 in S_2 is mapped upon the doubled C^3 in the fixed S_3 by means of the equations in (49).

3. The locus of the zero point in S_3 . In S_2 we find K as the locus of the zero point. Each curve of the system C^2 has one such point, which generates K by the variation of ω . Each curve of the system C^3 has on it the zero point. What is the equation of its locus? Since $z_i = t_i$ is the base point of the mapping system which maps C^2 upon C^3 , all the y 's vanish at this point, but as z approaches t the limiting position of the direction is that of the tangent to the polar conic C^2 at $z_i = t_i$. If the factors (ik) in (49) are replaced by the coördinates of the tangent to the polar conic at the point $z_i = t_i$, and if we set $z_i = t_i$ in the other factors, the y 's do not vanish, and become nonic functions of t which have a common factor $t_1 t_2 t_4$. However, a much simpler way to get this parametric representation of the locus of the zero point in S_3 is to solve the bilinearforms $\phi'_{01} = \phi'_{02} = \phi'_{04} = 0$ for y_i , and put $z_i = t_i$ in the result, from which the factor $t_1 t_2 t_4$ can be removed. These equations are:

$$\begin{aligned}
 (50) \quad y_0 &= - 14t_1^2 t_2^2 t_4^2, \\
 y_1 &= t_1^4 t_4^2 - 3t_1^3 t_2^3 - 5t_1 t_2^2 t_4^3, \\
 y_2 &= t_2^4 t_1^2 - 3t_2^3 t_4^3 - 5t_2 t_4^2 t_1^3, \\
 y_4 &= t_4^4 t_2^2 - 3t_4^3 t_1^3 - 5t_4 t_1^2 t_2^3.
 \end{aligned}$$

These equations map the locus of the zero point in S_2 upon a locus in the space of the y 's. The order of this locus is 18, for a plane section $(U'y) = 0$ gives a sextic in t which intersects K in 24 points, but we find that this variable sextic and K have 6 fixed intersections at the flex points $t_i = t_k = 0$ and consequently 18 variable ones. Hence the locus of the zero point in S_3 is a curve of order 18 and will be denoted by C^{18} .

It has already been pointed out that the order of the group of the y 's is double the order of the group of the z 's and that to express a form in y and z covariantly its points and planes in S_3 must appear squared. This C^{18} can evidently be represented covariantly if we take the 10 quadratic combinations of the y 's from the equations (50) from which we can eliminate the factor $t_1 t_2 t_4$ and thereby eliminate the fixed intersections each taken twice, and if we take in primed variables the corresponding quadratic combinations of the U 's as the coefficients of these quadratic combinations of the y 's. This form is

$$(51) \quad f(t'^3, t^9) = 0,$$

and is of the third order in t' , and of the ninth order in t . The number of variable points in which this nonic intersects K is 36, which is double the order of C^{18} , since its points appear squared in (51). If $t = t'$ in (51) we find a form of order 12 which is $K^3 + 16H^2$, where H is the Hessian of K . We can then say that the form (51) is the third polar of $K^3 + 16H^2$ plus covariant terms containing the line coordinates tt' . To obtain these further terms one would make use of the complete system of invariants and covariants of K which has been calculated and tabulated by Gordan.*

4. Summary. The results obtained may be briefly summarized. The well known elliptic modular functions associated with the elliptic norm curve E^7 and the algebraic relations connecting them have been readily found from the geometric point of view. The system of contact cubics in (37), the coordinates of the modular lines L_m and L'_m and the parametric representation of the locus of the zero point in S_3 are new types of functions. The system of modular functions $B\gamma$ (in Klein's notation)† which define a curve of order 14 has not been found.

If a pair of points in the involution on the curve E^7 is isolated, the quadratic irrationality associated with the curve E^7 is obtained. This irrationality can be obtained from the system y_i in (44). If we substitute the values of these y_i in any of the quadrics (19) (except those bilinear in y and z), ρ is obtained as the square root of the reciprocal of a conic $g(t^6, z^2)$. This conic has the form

$$(52) \quad g(t^6, z^2) = z_1^2(2t_1 t_2^4 t_4 + t_1^2 t_2 t_4^3) + z_2^2(2t_1 t_2 t_4^4) + z_4^2(t_1^4 t_2 t_4) + z_1 z_2(t_1^2 t_2^3 t_4 - 2t_1^3 t_4^3) + z_2 z_4(-4t_1^3 t_2^3 - t_2^5 t_4) - z_1 z_4(3t_2^3 t_4^3),$$

* *Mathematische Annalen*, vol. 17 (1880), pp. 217, 359.

† See K. F., vol. 2, p. 396-397.

and constitutes the part in z of a quadric on the curve E^7 whose part in y is the square of the half period plane (47).* We can now write down the parametric representation of the curve E^7 . It is

$$\begin{aligned}
 (53) \quad y'_0 &= + 2t_1^2 t_2^3 z_2 \quad (41) - (2t_1^2 t_2^2 t_4 z_2 + 2t_1 t_2^3 t_4 z_1) \quad (12), \\
 y'_1 &= - t_1 t_2^3 t_4 z_4 \quad (12) + (t_1 t_2^3 t_4 z_2 - t_1^3 t_2^2 z_1) \quad (41), \\
 y'_2 &= - t_2 t_4^3 t_1 z_1 \quad (24) + (t_2 t_4^3 t_1 z_4 - t_2^3 t_4^2 z_2) \quad (12), \\
 y'_4 &= - t_4 t_1^3 t_2 z_2 \quad (41) + (t_4 t_1^3 t_2 z_1 - t_4^3 t_1^2 z_4) \quad (24), \\
 z'_1 &= z_1 t_1 t_2 t_4 \sqrt{g(t^6, z^2)}, \\
 z'_2 &= z_2 t_1 t_2 t_4 \sqrt{g(t^6, z^2)}, \\
 z'_4 &= z_4 t_1 t_2 t_4 \sqrt{g(t^6, z^2)}.
 \end{aligned}$$

If t is on K the above system maps the doubled C^2 upon the E^7 . It should be noted that the y 's vanish for $z_i = t_i$ and the z 's vanish when z is on a half period point.†

* Professor Sharpe of Cornell pointed out this fact to me, as well as a method of eliminating an extraneous factor $t_1 t_2 t_4$ from the parametric representation of the curve E^7 . I append the method in a foot note at the end of the paper.

† All the terms in y'_i contain the factor $t_1 t_2 t_4$ except one term in y'_0 and this term contains a z_4 . If we now find the intersection of the pencil of lines through the point t , $\lambda_1(42) + \lambda_2(14) = 0$ and the polar conic C^2 , we get the following values for z :

$$\begin{aligned}
 z_1 &= \lambda_1^2(-2t_1^2 t_2 - t_4^3) + \lambda_2^2 t_1 t_2 t_4 - 3t_2^2 + t_4 \lambda_1 \lambda_2, \\
 z_2 &= \lambda_1^2 t_1 t_2^2 + \lambda_2^2(-2t_2^2 t_4 - t_1^3) - (2t_1^2 t_2 + t_4^3) \lambda_1 \lambda_2, \\
 z_4 &= \lambda_1^2 t_1 t_2 t_4 + \lambda_2^2 t_2 t_4^2 t_1^2 + t_4 \lambda_1 \lambda_2.
 \end{aligned}$$

Hence when these values are put in (53) the factor $t_1 t_2 t_4$ can be removed.

UNIVERSITY OF ILLINOIS,
URBANA, ILL.

VITA

Born at Mayo, Ky., August 10, 1889, son of Thomas Clinton and Margaret Wheeler-Woods. Received his elementary education in the public school in Hopewell District, Mercer County, Ky. Entered the academy connected with Georgetown College at Georgetown, Ky., in 1908 and graduated from Georgetown College in 1914 with A.B. degree. Spent the summer of 1914 as a graduate student in the University of Chicago. Held an Assistantship in Mathematics in the University of Maine during the year 1914-15, Instructor, 1915-17. Received A.M. degree from the University of Maine in 1916. Was an Assistant in Mathematics in the University of Illinois during the years 1917-20.

UNIVERSITY OF CALIFORNIA LIBRARY,
BERKELEY

**THIS BOOK IS DUE ON THE LAST DATE
STAMPED BELOW**

Books not returned on time are subject to a fine of 50c per volume after the third day overdue, increasing to \$1.00 per volume after the sixth day. Books not in demand may be renewed if application is made before expiration of loan period.

SEP 12 1924

Binder
Gaylord Bros.
Makers
Syracuse, N. Y.
PAT. JAN 21, 1903

523611 QM 245
190

UNIVERSITY OF CALIFORNIA LIBRARY

